Predicting Developmental Delays in Children: The Malaria Borth Cohort Study

Data Science Workshop

James Osei-Mensa

Thursday, March 20, 2025





BACKGROUND

- Developmental delays in children refer to when a child does not reach their developmental milestones at the expected times.
- WHO estimates that 8% of all children under 5 years have some type of developmental deficit (Global Research on Developmental Disabilities Collaborators 2018)
- Bello et al found 44.6% of 330 children with delays in various domains (Bello, Quartey, and Appiah 2013)
- Developmental milestones not checked routinely
- Early intervention has high impact



• **Cognitive Development**: Delays in thinking, learning, and problem-solving skills.



- Cognitive Development: Delays in thinking, learning, and problem-solving skills.
- **Physical Development**: Delays in gross motor skills (e.g., crawling, walking) and fine motor skills (e.g., grasping, drawing).



- Cognitive Development: Delays in thinking, learning, and problem-solving skills.
- Physical Development: Delays in gross motor skills (e.g., crawling, walking) and fine motor skills (e.g., grasping, drawing).
- **Communication Development**: Delays in speech and language skills, including understanding and using language.



- Cognitive Development: Delays in thinking, learning, and problem-solving skills.
- Physical Development: Delays in gross motor skills (e.g., crawling, walking) and fine motor skills (e.g., grasping, drawing).
- Communication Development: Delays in speech and language skills, including understanding and using language.
- **Social and Emotional Development**: Delays in interacting with others, forming relationships, and managing emotions.



• Genetic Factors: Down syndrome, Fragile X syndrome, etc



- Genetic Factors: Down syndrome, Fragile X syndrome, etc
- Prenatal Factors: Gestational age, birth weight, exposure to alcohol, drugs, or infections during pregnancy, poor maternal nutrition, and complications during birth



- Genetic Factors: Down syndrome, Fragile X syndrome, etc
- Prenatal Factors: Gestational age, birth weight, exposure to alcohol, drugs, or infections during pregnancy, poor maternal nutrition, and complications during birth
- **Environmental Factors**: Exposure to toxins, lack of stimulation, and inadequate nutrition



- Genetic Factors: Down syndrome, Fragile X syndrome, etc
- Prenatal Factors: Gestational age, birth weight, exposure to alcohol, drugs, or infections during pregnancy, poor maternal nutrition, and complications during birth
- **Environmental Factors**: Exposure to toxins, lack of stimulation, and inadequate nutrition
- Medical Conditions: Chronic illnesses, infections



- Genetic Factors: Down syndrome, Fragile X syndrome, etc
- Prenatal Factors: Gestational age, birth weight, exposure to alcohol, drugs, or infections during pregnancy, poor maternal nutrition, and complications during birth
- **Environmental Factors**: Exposure to toxins, lack of stimulation, and inadequate nutrition
- Medical Conditions: Chronic illnesses, infections
- Socioeconomic Factors: Poverty, lack of access to healthcare



- Genetic Factors: Down syndrome, Fragile X syndrome, etc
- Prenatal Factors: Gestational age, birth weight, exposure to alcohol, drugs, or infections during pregnancy, poor maternal nutrition, and complications during birth
- Environmental Factors: Exposure to toxins, lack of stimulation, and inadequate nutrition
- Medical Conditions: Chronic illnesses, infections
- Socioeconomic Factors: Poverty, lack of access to healthcare
- Parental Factors: Parental mental health issues, lack of parental education



SURVEY INSTRUMENT

- Developmental Milestones Checklist III
- Domains surveyed
 - Gross motor
 - ▶ Fine motor
 - Language
- Children examined at yearly intervals from 12 months to 60 months




MODEL EVALUATION

Table 1: Performance metrics of different models. MSE refers to Mean Squared Error, and RMSE refers to Root Mean Squared Error.

Model	MSE (Train)	MSE (Test)	RMSE (Train)	RMSE (Test)
Ridge Regression	45.6690	51.7989	6.7537	7.1781
Support Vector Regression	25.6013	53.2237	5.0515	7.2675
Random Forest	6.8531	50.3259	2.6161	7.0750
Decision Tree	0	51.3561	0	7.1469
Multi Layer Perceptron	30.8844	50.6152	5.5532	7.0930



IMPORTANT PREDICTORS



Sex Walk at 1 year Stand at 1 year MUAC at 1 year (cm) Apgar at 5 min Birth weight (kg) Feature value Birth head circumference (cm) Birth length (cm) Multiple birth Length for age z-score at 1 year Mother's age (years) Gestational age (weeks) Gynaecological Abnormally SHAP Value (Impact on Predicted Developmental Score)

Figure 1: Key Clinical and Growth Factors Affecting Child Development

Figure 2: Impact of Clinical and Growth Factors on Developmental Scores

LIMITATIONS

• Sampled from only one population area



LIMITATIONS

- Sampled from only one population area
- Small sample size



LIMITATIONS

- Sampled from only one population area
- Small sample size
- Large gaps between survey times



CONCLUSION

- Prediction model based on only 5 easily obatinable variables
- Variables in final model: Sex, Walking at 1 year, Standing at 1 year, Muac score at 1 year, Apgar score at 5 minutes
- The model can predict developmental score at an error margin of 6.9327
- The model, however, needs to be validated in a different cohort



REFERENCES

Bello, Ajediran I., Jonathan NA Quartey, and Louisa A. Appiah. 2013. "Screening for Developmental Delay Among Children Attending a Rural Community Welfare Clinic in Ghana." *BMC Pediatrics* 13 (1): 119. https://doi.org/10.1186/1471-2431-13-119.

Global Research on Developmental Disabilities Collaborators. 2018. "Developmental Disabilities Among Children Younger Than 5 Years in 195 Countries and Territories, 1990–2016: A Systematic Analysis for the Global Burden of Disease Study 2016 - ClinicalKey." Lancet Global Health 6 (10): e1100–1121.

